



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Using the pimeloyl-CoA synthetase adenylation fold to synthesize fatty acid thioesters

**Citation for published version:**

Wang, M, Moynié, L, Harrison, PJ, Kelly, V, Piper, A, Naismith, JH & Campopiano, DJ 2017, 'Using the pimeloyl-CoA synthetase adenylation fold to synthesize fatty acid thioesters', *Nature Chemical Biology*, vol. 13, pp. 660–667. <https://doi.org/10.1038/nchembio.2361>

**Digital Object Identifier (DOI):**

[10.1038/nchembio.2361](https://doi.org/10.1038/nchembio.2361)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Other version

**Published In:**

Nature Chemical Biology

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Figure 1





Figure 2




Figure 3




Figure 4




Figure 5

